

Diaphragm

? Diaphragm

Introduction

- The **diaphragm** is a **thin, dome-shaped musculetendinous partition** separating the **thoracic** and **abdominal cavities**.
 - It is the **chief muscle of respiration** — contraction increases thoracic volume, causing inspiration.
 - It is convex upward (right dome higher due to liver).
 - At rest, its highest point (right dome) reaches the level of the **5th rib**.
 - It descends up to **one intercostal space** during deep inspiration.
-

Gross Anatomy

The diaphragm consists of:

1. **Peripheral Muscular Part**
 2. **Central Tendinous Part**
-

1. Muscular Part

Arises from three main parts:

a. Sternal Part

- From **posterior surface of xiphoid process** (two small slips).
- Separated by small gap — *for superior epigastric vessels*.

b. Costal Part

- From **inner surfaces of lower six ribs and their costal cartilages** (interdigitating with transversus abdominis).
- Forms largest portion.

c. Vertebral Part

- Arises by **crura and arcuate ligaments**:
 - **Right crus**: from bodies of L1–L3 and intervertebral discs.
 - **Left crus**: from bodies of L1–L2.
 - The **right crus** is longer.
- The two crura are connected by **median arcuate ligament** (in front of aorta).

Arcuate Ligaments:

- **Median arcuate ligament** ? over aortic opening.
 - **Medial arcuate ligament** ? thickened fascia over psoas major.
 - **Lateral arcuate ligament** ? thickened fascia over quadratus lumborum.
-

2. Central Tendon

- Strong, trifoliate (three-leafed) aponeurotic sheet.
 - Lies opposite **xiphoid process (T8–T9 level)**.
 - The **pericardium** is fused to its central portion.
-

Openings in the Diaphragm

There are **three large openings** and several small ones.

1. Caval Opening (T8)

- Lies in **central tendon (right leaf)**.
 - **Structures passing:**
 - Inferior vena cava.
 - Terminal branches of right phrenic nerve.
 - **Mnemonic: “V8” — Vena cava at T8.**
 - Function: enlarges during inspiration ? facilitates venous return.
-

2. Oesophageal Opening (T10)

- Lies in **muscular part (right crus)**.
- **Structures passing:**

- Oesophagus.
 - Vagal trunks (anterior and posterior).
 - Oesophageal branches of left gastric vessels.
 - **Mnemonic: “Oesophagus Ten (10)”.**
 - Function: acts as a sphincter — diaphragm assists in preventing reflux.
-

3. Aortic Opening (T12)

- Lies **behind the median arcuate ligament**.
 - **Structures passing:**
 - Aorta.
 - Thoracic duct.
 - Azygos vein.
 - **Mnemonic: “Aortic Twelve (12)”.**
 - Function: not affected by diaphragmatic contraction (lies posterior).
-

Other Small Openings

- **Right crus:** greater and lesser splanchnic nerves.
- **Left crus:** hemiazygos vein.

- **Between crura:** sympathetic trunk.
 - **Sternocostal triangle:** superior epigastric vessels.
 - **Behind lateral arcuate ligament:** subcostal vessels and nerve.
-

Relations

Superior (Thoracic) Surface

Covered by pleura and pericardium.

- **Right dome:** related to liver and right lung base.
- **Left dome:** related to stomach, spleen, and left lung base.

Inferior (Abdominal) Surface

Covered by peritoneum, related to:

- Right dome ? right lobe of liver, right kidney, suprarenal gland.
 - Left dome ? left lobe of liver, fundus of stomach, spleen, left kidney, suprarenal gland.
 - Central tendon ? pericardium (via fibrous layer).
-

Important Vertebral Levels (Mnemonics)

- **T8 ? Vena Cava (V8)**
 - **T10 ? Oesophagus (OesophaTen)**
 - **T12 ? Aorta (Aortwelve)**
-

? Nerve Supply

1. Motor Nerve

- **Phrenic nerve (C3, C4, C5)** — “**C3, 4, 5 keep the diaphragm alive.**”
- Right phrenic nerve supplies the **right half**, and the left phrenic nerve supplies the **left half**.
- The central tendon region of the diaphragm is entirely **phrenic-supplied**.

2. Sensory Nerves

- **Central part:** Phrenic nerve (pain referred to shoulder – C4 dermatome).
- **Peripheral part:** Lower **six intercostal nerves** and **subcostal nerve (T12)**.
- Pain from these peripheral parts is referred to the **costal margins or abdominal wall**.

? Actions of the Diaphragm

1. Chief Muscle of Inspiration

- On contraction, domes descend ? thoracic cavity enlarges ? pressure drops ? air enters lungs.

2. Expiration (Relaxation Phase)

- When the diaphragm relaxes, domes rise ? thoracic volume decreases ? air expelled.

3. Assists in Circulation

- Contraction decreases intrathoracic pressure and increases intra-abdominal pressure, aiding **venous return** via IVC and **lymph flow** via thoracic duct.

4. Acts as a Partition and Sphincter

- Through its openings, the diaphragm controls passage of oesophagus, aorta, IVC, etc.
- The **right crus acts as a physiological sphincter** for the lower oesophagus.

5. Abdominal Pressurization

- With other muscles, helps in **micturition, defecation, parturition, and vomiting** by raising intra-abdominal pressure.

? Dissection Highlights

- The diaphragm is seen from the abdominal cavity after removing abdominal viscera.
- Identify:
 - **Central tendon** – shiny and aponeurotic.
 - **Crura** – muscular slips attached to lumbar vertebrae.
 - **Arcuate ligaments** – median, medial, and lateral.

- **Openings** – IVC at T8 (in tendon), oesophagus at T10 (in muscle), aorta at T12 (behind ligament).
 - Observe the **phrenic nerves** descending anteriorly and **sympathetic trunks** posteriorly.
-

?? Clinical Anatomy

1. Paralysis of Diaphragm

- Due to **phrenic nerve injury** (e.g., neck trauma, mediastinal tumor).
- On X-ray ? **paradoxical movement** (dome moves upward during inspiration).

2. Eventration of Diaphragm

- Congenital thinning or absence of muscle ? one dome balloons upward.
- Commonly on left side; may compress lung.

3. Hiatus Hernia

- Herniation of stomach into thorax through the **oesophageal opening**.
 - *Sliding type*: gastroesophageal junction moves upward.
 - *Paraesophageal type*: fundus herniates beside oesophagus.

4. Congenital Diaphragmatic Hernia (Bochdalek type)

- Due to failure of closure of **pleuroperitoneal canal** (posterolateral defect).

- Abdominal viscera herniate into thorax ? pulmonary hypoplasia in newborn.

5. Morgagni Hernia

- Herniation through **sternocostal triangle (foramen of Morgagni)** – rare, anterior type.

6. Phrenic Pain Referred to Shoulder

- Central diaphragmatic pain ? referred to **shoulder (C4)** via phrenic nerve.
- Peripheral pain ? referred to **costal margins** (intercostal nerves).

7. Subphrenic Abscess

- Pus collects below diaphragm (usually right side) following peritonitis or liver abscess.
- Causes elevation of diaphragm and referred shoulder pain.

8. Accessory Phrenic Nerve

- May arise from **nerve to subclavius (C5)** ? joins phrenic nerve in thorax.
- Damage to both may cause partial paralysis.

? Development of Diaphragm

The diaphragm develops from **four embryonic components**:

1. Septum Transversum

- Forms **central tendon**.

- Lies opposite cervical somites (C3–C5) ? hence phrenic nerve origin.

2. **Pleuroperitoneal Membranes**

- Close the **pleuroperitoneal canals** on each side posteriorly.

3. **Dorsal Mesentery of Oesophagus**

- Forms **crura** and **median portion** around oesophageal opening.

4. **Body Wall (Peripheral Muscular Part)**

- Contributes to **costal portion** of diaphragm.

Timeline:

- Fusion complete by the **8th week** of development.
- Initially cervical in position, descends to thoracic level by 8th week, dragging phrenic nerve downward — explaining its long course.

Clinical Correlation:

- Failure of fusion ? **congenital diaphragmatic hernia** (Bochdalek type).
- Defect in septum transversum ? **eventration** or **Morgagni hernia**.

? Mnemonics – Diaphragm

1. Openings of Diaphragm (Levels and Structures)

Mnemonic: ? “I 8 10 Eggs At 12”

- **I 8 ? IVC at T8 ?** Inferior Vena Cava + Right Phrenic Nerve.
 - **10 Eggs ? Oesophagus at T10 ?** Oesophagus + Vagal Trunks + Left Gastric Vessels.
 - **At 12 ? Aorta at T12 ?** Aorta + Thoracic Duct + Azygos Vein.
-

2. Structures Passing Through Crura

Mnemonic: “Silly Great Lads Hide Behind Crura”

- **S** – Sympathetic trunks.
 - **G** – Greater splanchnic nerves.
 - **L** – Lesser splanchnic nerves.
 - **H** – Hemiazygos vein.
 - **B** – Both crura connect via Median Arcuate Ligament.
-

3. Arcuate Ligaments (Attachments)

Mnemonic: “L-M-M: Laterally Quadratus, Medially Psoas, Middle Aorta.”

- **Lateral arcuate ligament ?** over **Quadratus lumborum**.
 - **Medial arcuate ligament ?** over **Psoas major**.
 - **Median arcuate ligament ?** arches over **aorta**.
-

4. Functions of Diaphragm

Mnemonic: “AIR PRESS”

A – Aids in **Abdominal pressure** (urination, defecation, parturition).

I – **Inspiration** (main function).

R – **Returns venous blood** to heart.

P – **Partition** between thorax and abdomen.

R – **Reflex control** of breathing via phrenic.

E – **Expulsion** (vomiting, cough).

S – **Sphincter action** at oesophageal opening.

S – **Support** to abdominal viscera.

5. Components of Diaphragm Development

Mnemonic: “Several Parts Develop Diaphragm.”

- **S** – Septum transversum ? central tendon.
 - **P** – Pleuroperitoneal membranes.
 - **D** – Dorsal mesentery of oesophagus.
 - **D** – Derived muscle from body wall (peripheral part).
-

6. Nerve Supply

Mnemonic: “C3, 4, 5 keep the diaphragm alive.”

– Phrenic nerve origin from C3–C5 cervical segments.

? Facts to Remember – Diaphragm

- The **diaphragm** is the **principal muscle of inspiration**.
 - It separates the **thoracic and abdominal cavities**.
-

- Formed of **peripheral muscle fibers** and a **central tendon**.
- The **right dome** lies higher (due to liver) than the **left**.
- **Central tendon** lies at the level of **xiphoid process (T8–T9)**.
- **Three major openings:**
 - **IVC – T8** (in tendon).
 - **Oesophagus – T10** (in right crus).
 - **Aorta – T12** (behind median arcuate ligament).
- **Arcuate ligaments:** median (over aorta), medial (over psoas), lateral (over quadratus lumborum).
- **Right crus** is longer and wider than the left.
- **Nerve supply:**
 - Motor ? Phrenic nerve (C3–C5).
 - Sensory ? Phrenic (central part) and lower 6 intercostal + subcostal (peripheral part).
- **Blood supply:** Musculophrenic, pericardiophrenic, superior and inferior phrenic arteries.
- **Action:** Contraction ? inspiration by lowering domes; relaxation ? expiration.
- During inspiration, the **central tendon descends about 1–2 cm** in quiet breathing, up to 5 cm in deep breathing.

- **Paralysis** of one half ? elevated dome on that side with paradoxical movement.
- **Eventration:** Congenital deficiency causing ballooning of diaphragm.
- **Hiatus hernia:** Protrusion of stomach through oesophageal opening.
- **Congenital diaphragmatic hernia:** Failure of pleuroperitoneal membrane closure (Bochdalek type).
- **Phrenic pain** is referred to the **shoulder (C4)**; peripheral pain to costal margins.
- The diaphragm develops from four embryological sources: septum transversum, pleuroperitoneal membranes, dorsal mesentery of oesophagus, and body wall.
- **Central tendon** derived from septum transversum.
- **Clinical importance:** Vital in respiration and venous return to heart.

?? Clinicoanatomical Problems – Diaphragm

1. Case of Diaphragmatic Paralysis

A 42-year-old man presents with breathlessness. X-ray shows the right dome of diaphragm elevated and moving paradoxically on inspiration.

? **Diagnosis:** Paralysis of diaphragm due to right **phrenic nerve injury** (C3–C5).

2. Case of Shoulder Tip Pain After Liver Abscess

A patient with right subphrenic abscess complains of pain over right shoulder tip.

? **Explanation:** Referred pain from diaphragm via **phrenic nerve (C4 dermatome)**.

3. Eventration of Diaphragm

A child has abnormally high position of the left dome of diaphragm on X-ray, but no rupture.

? **Diagnosis: Eventration** — congenital thinning or partial absence of muscle fibers, causing ballooning.

4. Congenital Diaphragmatic Hernia (Bochdalek Type)

A newborn has severe respiratory distress immediately after birth. X-ray shows bowel loops in the thorax.

? **Diagnosis: Posterolateral diaphragmatic hernia** due to failure of **pleuroperitoneal canal closure**.

5. Morgagni Hernia

An elderly woman with chronic cough presents with epigastric bulge near xiphoid process.

? **Cause:** Herniation through **sternocostal triangle (foramen of Morgagni)**.

6. Sliding Hiatus Hernia

Middle-aged patient with heartburn and regurgitation. Barium swallow shows stomach and gastroesophageal junction above diaphragm.

? **Diagnosis: Sliding hiatus hernia** through oesophageal opening (T10).

7. Paraesophageal Hiatus Hernia

A patient's stomach fundus is seen herniating beside oesophagus without displacement of gastroesophageal junction.

? **Diagnosis: Paraesophageal hernia** – dangerous due to possible strangulation.

8. Referred Epigastric Pain

Pain perceived in epigastrium when diaphragmatic pleura irritated (e.g. subphrenic abscess).

? **Reason:** Central diaphragmatic part supplied by **phrenic nerve**, referred to epigastrium.

9. Bilateral Diaphragmatic Paralysis

After cervical spinal cord injury at C4 level, patient develops respiratory failure.

? **Reason: Phrenic nerve roots (C3–C5)** are affected, diaphragm paralyzed ? ventilatory failure.

10. Post-operative Respiratory Difficulty

After open heart surgery, patient has raised left dome on X-ray.

? **Cause:** Iatrogenic injury to **left phrenic nerve** (lies on pericardium).

11. Phrenic Nerve Palsy During Neck Surgery

Patient develops dyspnea after thyroidectomy.

? **Reason:** Accidental ligation of **phrenic nerve** crossing anterior to scalenus anterior muscle.

12. Congenital Absence of Diaphragm

Neonate with respiratory distress has no separation between thoracic and abdominal cavities.

? **Diagnosis: Agenesis of diaphragm** – failure of all four embryonic components to fuse.

13. Accessory Phrenic Nerve Damage

Partial diaphragmatic paralysis occurs though main phrenic nerve preserved.

? **Explanation:** Injury to **accessory phrenic nerve** (from nerve to subclavius, C5).

14. Hiccups (Singultus)

Persistent involuntary contraction of diaphragm causing sudden inspiration with glottis closure.

? **Reason:** Reflex irritation of **phrenic nerve or medullary centers**.

15. Subphrenic Abscess

Post-appendectomy patient develops fever and shoulder pain.

? **Diagnosis: Subphrenic abscess** beneath diaphragm due to pus collection.

? **Feature:** Elevation of diaphragm and referred shoulder pain (C4).

16. Abnormal Movement in Fluoroscopy

During inspiration, one dome moves downward but the other moves upward.

? **Diagnosis: Paradoxical movement** – sign of unilateral diaphragmatic paralysis.

17. Diaphragmatic Rupture After Trauma

A man sustains blunt abdominal trauma; stomach loops visible in left thoracic cavity.

? **Diagnosis: Traumatic rupture of diaphragm** – more common on left side (liver protects right).

18. Congenital Eventration Mistaken for Hernia

Infant's X-ray shows high dome with gut shadow beneath intact pleura.

? **Differentiation:** Eventration (no rupture) vs hernia (defect present).

19. Diaphragmatic Hernia in Adults

An adult with chronic cough develops herniation of omentum through **foramen of Bochdalek** (posterolateral).

? **Significance:** Late presentation of congenital defect.

20. Phrenic Nerve Involvement in Pericarditis

Patient with pericardial irritation develops pain radiating to shoulder tip.

? **Reason:** Common sensory pathway via **phrenic nerve** (C4).

? Frequently Asked Questions — Diaphragm

1. Describe the origin, insertion, nerve supply, and actions of the thoracoabdominal diaphragm.
2. Write short notes on:

- Major openings of diaphragm and the structures passing through them.
- Minor openings and their contents.
- Diaphragmatic hernia — congenital and acquired types.
- Course and distribution of the phrenic nerve.

3. **Explain why the cardiac end of the stomach is prone to sliding hernia.**

- Due to weakness of the **phrenico-oesophageal membrane**, allowing the cardiac end to slide through the oesophageal hiatus, disturbing the **valvular mechanism** at the gastroesophageal junction and causing **reflux**.

4. **List the vertebral levels of major openings of the diaphragm and structures transmitted through each.**

- T8 ? Inferior vena cava, right phrenic nerve.
- T10 ? Oesophagus, vagal trunks, left gastric vessels.
- T12 ? Aorta, thoracic duct, azygos vein.

5. **Discuss congenital anomalies of diaphragm development.**

- Posterolateral (Bochdalek) hernia, Retrosternal (Morgagni) hernia, Eventration, and Complete absence of diaphragm.

6. **What is the nerve supply of the diaphragm?**

- **Motor:** Phrenic nerve (C3–C5).

- **Sensory:** Central part by phrenic nerve; peripheral part by lower six intercostal nerves.

? Multiple Choice Questions — Diaphragm

1. Which of the following structures does *not* pass through the diaphragm?

- a. Oesophagus
- b. Aorta
- c. **Cisterna chyli ?**
- d. Inferior vena cava

To build a complete **exam-ready MCQ section**, here are **additional standard questions** (patterned on BD Chaurasia and AIIMS/NEET PG level):

2. The diaphragm is supplied by which nerve?

- a. Intercostal nerve
- b. Vagus nerve
- c. **Phrenic nerve ?**
- d. Sympathetic nerve

3. The oesophagus passes through the diaphragm at the level of:

- a. T8
- b. **T10 ?**
- c. T12
- d. L1

4. The inferior vena cava passes through diaphragm at the level of:

- a. **T8 ?**
 - b. T10
 - c. T12
 - d. T6
-

5. The aortic opening of diaphragm lies at:

- a. T10
 - b. **T12 ?**
 - c. L1
 - d. T9
-

6. The phrenic nerve arises from:

- a. C2–C3
 - b. C4–C6
 - c. **C3–C5 ?**
 - d. T1–T3
-

7. Which of the following does not pass through oesophageal opening?

- a. Oesophagus
 - b. Vagus nerve
 - c. Left gastric artery
 - d. **Azygos vein ?**
-

8. The central tendon of diaphragm is derived from:

- a. **Septum transversum ?**
 - b. Pleuroperitoneal membrane
 - c. Dorsal mesentery of oesophagus
 - d. Lateral body wall
-

9. Which dome of diaphragm is higher?

- a. **Right ?**
 - b. Left
 - c. Equal
 - d. Variable
-

10. The muscle fibers of diaphragm insert into:

- a. Costal arch
 - b. **Central tendon ?**
 - c. Vertebral column
 - d. Linea alba
-

11. Which structure is behind the diaphragm?

- a. Aorta
 - b. **Thoracic duct ?**
 - c. Inferior vena cava
 - d. Oesophagus
-

12. Eventration of diaphragm is due to:

- a. Trauma
 - b. Inflammation
 - c. **Congenital muscle defect ?**
 - d. Hernia
-

13. Which artery does *not* supply the diaphragm?

- a. Pericardiophrenic
 - b. Musculophrenic
 - c. **Inferior mesenteric ?**
 - d. Inferior phrenic
-

14. The main muscle of inspiration is:

- a. External intercostals
 - b. Internal intercostals
 - c. **Diaphragm ?**
 - d. Scalene
-

15. The right crus of diaphragm arises from:

- a. L1–L2
 - b. **L1–L3 ?**
 - c. L2–L4
 - d. T12–L2
-

16. The left crus of diaphragm arises from:

- a. L1–L3
 - b. **L1–L2 ?**
 - c. L2–L3
 - d. T12–L2
-

17. Pain from central part of diaphragm is referred to:

- a. Epigastrium
 - b. **Shoulder (C4 dermatome) ?**
 - c. Lower chest
 - d. Back
-

18. Posterolateral diaphragmatic hernia is also called:

- a. Morgagni hernia
 - b. **Bochdalek hernia ?**
 - c. Hiatus hernia
 - d. Paraesophageal hernia
-

19. Which structure passes behind the median arcuate ligament?

- a. Inferior vena cava
 - b. Oesophagus
 - c. **Aorta ?**
 - d. Vagus nerve
-

20. All the following are openings in the diaphragm except:

- a. Caval opening
- b. Oesophageal hiatus
- c. Aortic hiatus
- d. **Renal opening ?**

? Viva Voce — Diaphragm

1. What is the diaphragm?

A dome-shaped **musculotendinous partition** separating the **thoracic and abdominal cavities**; the **chief muscle of inspiration**.

2. What are the parts of the diaphragm?

- **Muscular part** – sternal, costal, and vertebral.
 - **Central tendinous part** – aponeurotic and trifoliate in shape.
-

3. Which dome of diaphragm is higher?

The **right dome**, because it rests on the liver.

4. What is the nerve supply of the diaphragm?

- **Motor:** Phrenic nerve (C3, C4, C5).
 - **Sensory:** Phrenic (central part) and lower six intercostal + subcostal nerves (peripheral part).
-

5. What is the mnemonic for openings of the diaphragm?

“I 8 10 Eggs At 12”

- IVC ? T8
 - Oesophagus ? T10
 - Aorta ? T12
-

6. Which opening lies in the central tendon?

Caval opening (T8) for inferior vena cava and right phrenic nerve.

7. What structures pass through the oesophageal opening?

- Oesophagus
 - Anterior and posterior vagal trunks
 - Oesophageal branches of left gastric vessels.
-

8. What structures pass through the aortic opening?

- Aorta
 - Thoracic duct
-

- Azygos vein.
-

9. Which crus of diaphragm is longer?

The **right crus** (arises from L1–L3) is longer than the left (L1–L2).

10. What is the function of the crura?

They form a **slings** around the oesophagus and act as a **physiological sphincter** preventing gastric reflux.

11. Name the three arcuate ligaments.

- **Median arcuate ligament** – arches over aorta.
 - **Medial arcuate ligament** – over psoas major.
 - **Lateral arcuate ligament** – over quadratus lumborum.
-

12. What are the actions of the diaphragm?

- **Inspiration:** Increases thoracic volume by descending.
 - **Expiration:** Relaxes and ascends.
 - **Abdominal compression:** Assists in defecation, micturition, parturition.
 - **Venous return:** Aids blood flow via IVC.
-

13. What is the developmental origin of diaphragm?

From **four embryonic sources**:

1. Septum transversum
 2. Pleuroperitoneal membranes
 3. Dorsal mesentery of oesophagus
 4. Muscular ingrowths from body wall.
-

14. What is the central tendon derived from?

Septum transversum.

15. What are the common congenital anomalies of diaphragm?

- **Bochdalek hernia** (posterolateral)
 - **Morgagni hernia** (sternocostal)
 - **Eventration** (muscle defect)
 - **Complete absence** (rare).
-

16. What is the vertebral level of the diaphragm's central tendon?

Approximately **T8–T9**.

17. What is eventration of diaphragm?

A congenital condition where one dome is elevated due to **defective muscular development**, replaced by thin membrane.

18. What is hiatus hernia?

Herniation of the **stomach** through the **oesophageal hiatus** into thorax; may be **sliding** or **paraesophageal**.

19. What is referred pain of diaphragm and why?

Pain from the diaphragm is felt at the **shoulder tip (C4)** because both **phrenic** and **supraclavicular nerves** share the same spinal root value.

20. What is the difference between Bochdalek and Morgagni hernias?

- **Bochdalek:** Posterolateral defect (common).
 - **Morgagni:** Anterior retrosternal defect (rare).
-

21. What is paradoxical movement of diaphragm?

In unilateral diaphragmatic paralysis, the paralyzed dome moves **upward during inspiration** instead of descending.

22. Why is right phrenic nerve important surgically?

It's short and runs over the **inferior vena cava**, hence easily damaged during hepatic or pericardial surgery.

23. What is the blood supply of diaphragm?

- **Arteries:** Musculophrenic, pericardiophrenic, superior & inferior phrenic arteries.
 - **Veins:** Drain into azygos, inferior phrenic, and internal thoracic veins.
-

24. What are the functions of the diaphragm other than respiration?

Aids in **vomiting, micturition, defecation, parturition**, and **cough reflex**.

25. What is the Organ of Bochdalek?

Not an organ — refers to **Bochdalek's foramen**, the site of **posterolateral congenital diaphragmatic hernia**.

26. Which muscle forms the floor of thoracic cavity?

Diaphragm.

27. What happens if the phrenic nerve is cut?

Paralysis of diaphragm on that side ? elevation of dome and **loss of respiratory function**.

28. Why does the diaphragm descend during inspiration?

Because of **phrenic nerve stimulation** ? contraction ? increase in vertical thoracic diameter.

29. What is the function of the oesophageal hiatus in preventing reflux?

The **right crus** forms a sling around the oesophagus acting as an **external sphincter**.

30. Why is congenital diaphragmatic hernia dangerous in newborns?

It causes **compression of developing lungs**, leading to **pulmonary hypoplasia and respiratory distress**.